



Official reprint from UpToDate®

www.uptodate.com ©2021 UpToDate, Inc. and/or its affiliates. All Rights Reserved.

Wolters Kluwer

Lesbian, gay, bisexual, and other sexual minoritized youth: Epidemiology and health concerns

Authors: Michelle Forcier, MD, MPH, Johanna Olson-Kennedy, MD**Section Editor:** Diane Blake, MD**Deputy Editor:** Mary M Torchia, MDAll topics are updated as new evidence becomes available and our [peer review process](#) is complete.**Literature review current through:** Apr 2021. | **This topic last updated:** Sep 17, 2020.

INTRODUCTION

The term "sexual minoritized" encompasses a variety of gender and sexual identities and expressions that differ from cultural norms (eg, lesbian, gay, bisexual, transgender), as well as identities and expressions that defy discrete labels [1].

This topic will focus on the epidemiology and health concerns of sexual minoritized youth who identify themselves as lesbian, gay, bisexual, pansexual, or unsure (questioning) of their sexual identity, who avoid discrete sexual orientation labels. An overview of primary care for such youth is presented separately. (See ["Lesbian, gay, bisexual, and other sexual minoritized youth: Primary care"](#).)

Gender diversity in children and adolescents is discussed separately. (See ["Gender development and clinical presentation of gender diversity in children and adolescents"](#) and ["Management of transgender and gender-diverse children and adolescents"](#).)

TERMINOLOGY

- **Sexual orientation** – Sexual orientation is an individual's pattern of physical and emotional arousal (including fantasies, activities, and behaviors) and the gender(s) of

persons to whom an individual is physically or sexually attracted [2].

- Homosexual sexual orientation refers to sexual attraction to same-sex individuals
 - Heterosexual sexual orientation refers to sexual attraction to opposite-sex individuals
 - Bisexual sexual orientation refers to sexual attraction to both same-sex and opposite-sex individuals
 - Pansexual sexual orientation refers to sexual attraction to individuals of any gender identity or biologic sex
 - Asexual sexual orientation refers to lack of sexual attraction to any individuals
- **Sexual identity** – Sexual identity is an individual's assessment of his or her sexual orientation; male individuals who self-identify as homosexual are often referred to as gay; female individuals who self-identify as homosexual are often referred to as lesbian [3]. Youth who self-identify as heterosexual are often referred to as straight. Youth who are unsure of or struggle with their sexual identity may be referred to as "questioning" [4].

The individual youth is the best person to describe and define his or her identity. Some adolescents prefer the terms "mostly heterosexual" or "mostly homosexual," to reflect their feeling of being between categories [5]. Others may avoid categorization, preferring more diffuse terminology such as queer, pansexual, or fluid regarding their sexuality.

- **Sexual behavior** – Sexual behavior refers to particular sexual activities and incorporates the gender(s) of sexual partners. Sexual attraction and identity correlate closely, but not completely, with reports of sexual behavior [6-11]. Youth who self-identify as heterosexual may engage in sexual activity with same-sex partners; youth who self-identify as homosexual may remain sexually inexperienced. Sexual behavior does not necessarily indicate sexual orientation; it may represent experimentation, exploration, or exploitation [12]. Sexual behaviors are more important than self-identified labels in assessing health risk. (See "[Lesbian, gay, bisexual, and other sexual minoritized youth: Primary care](#)", section on 'Sexuality'.)
- **Sexual minoritized** – The term "sexual minoritized" encompasses a variety of gender and sexual identities and expressions that differ from cultural norms (eg, lesbian, gay, bisexual, transgender), as well as identities and expressions that defy discrete labels

[1].

This topic will focus on sexual minoritized youth who identify themselves as lesbian, gay, bisexual, or unsure (questioning) of their sexual identity, who avoid discrete sexual orientation labels, and who have had sexual contact with persons of the same sex or persons of both sexes. Gender diversity is discussed separately. (See "[Gender development and clinical presentation of gender diversity in children and adolescents](#)" and "[Management of transgender and gender-diverse children and adolescents](#)".)

DEVELOPMENTAL PERSPECTIVE

Sexual orientation has multiple dimensions, including attractions, identity, behaviors, and the gender of sexual partners, all of which are important to understanding developing sexuality ([figure 1](#)) [3,13,14]. (See "[Adolescent sexuality](#)".)

Sexual exploration, experimentation, and discovery are part of the normal process of incorporating sexuality into one's sexual identity [15-17]. Before puberty, many children experiment with cross-gender play and expression. Cross-gender interests and expression in the prepubertal years are neither necessarily nor predictably associated with adolescent or adult sexual orientation. However, gender-diverse prepubertal children more often identify as gay or lesbian than transgender in adolescence and adulthood [18-21]. (See "[Gender development and clinical presentation of gender diversity in children and adolescents](#)", [section on 'Prepubertal children'](#).)

As children approach puberty and adolescence, sexuality (ie, attraction, interests, orientation, and behaviors) becomes relevant to the life tasks of developing relationships, intimacy, and creating family. Defining one's sexual orientation occurs over time. First awareness of same-sex attraction, which may or may not persist, has been described as early as age 9 for boys and 10 for girls, but sexual minoritized youth may self-identify orientation at later ages than heterosexual peers [22,23].

Formation of sexual identity among youth may be fluid [16,24]. Adolescents explore their emerging sexuality through attractions, fantasies, and behaviors [16,25]. Adolescent sexual identity and self-identified orientation does not necessarily correlate with sexual behaviors [6-11,26]. Adolescents may have same-sex attractions and fantasies but not

identify as homosexual; they may identify as homosexual but remain sexually inexperienced; or they may identify as heterosexual but engage in same sex behaviors. (See ["Adolescent sexuality", section on 'Adolescent development'](#).)

Strong family and societal expectations may influence the experience and expression of gender and sexuality in children and adolescents [27,28]. Some parents and families may try to redirect gender diverse expression and same-sex attractions, identity, or behavior to fit into heteronormative social parameters (ie, those that view opposite-sex relationships as the only "normal" relationships in society) [29]. Peer pressure may lead adolescents to behaviors that have little to do with attraction. Externalized and internalized sexual prejudice (also known as "homophobia" or "homonegativity") may lead adolescents to avoid sexual activity altogether, adopt heterosexual activity, or mistreat others who they perceive as homosexual [30]. (See ["Gender development and clinical presentation of gender diversity in children and adolescents", section on 'Gender development in childhood'](#).)

EPIDEMIOLOGY

Although epidemiologic studies of sexual minoritized youth use widely different measures [14], in several large surveys, 5 to 12 percent of high school youth report same-sex attractions or behaviors [31,32]. In a nationwide survey of high-school students, approximately 44 percent reported sexual contact with only the opposite sex, 2 percent with only the same sex, and 5 percent with both sexes [33]. Sexual minoritized students were racially, ethnically, socially, economically, and geographically diverse.

POTENTIAL PSYCHOSOCIAL AND HEALTH CONCERNS

Although the majority of sexual minoritized youth are healthy and well adjusted [17,34,35], some sexual minoritized youth are at increased risk for adverse psychosocial and health outcomes. Compared with heterosexual peers, sexual minoritized youth have increased rates of victimization, depression, suicide, substance use, homelessness, sexually transmitted infections, and unplanned pregnancy [32,36-43]. These health disparities are not intrinsic to sexual minoritized status but appear to be related to stigmatization and lack of acceptance of specific sexual behaviors, identities, and orientations [44-52].

Stigmatization and minority stress — Sexual minoritized youth may be stigmatized and marginalized by individuals or in communities and cultures that are dismissive, openly rejecting, or hostile [14]. Stigmatization is an important moderator of the disparities in psychosocial and health outcomes of sexual minoritized youth [44-46,49-51,53-56]. In surveys of sexual minoritized young adults, high levels of family rejection during adolescence was associated with suicidality, depression, illegal drug use, and engaging in unprotected sex; family acceptance was associated with greater self-esteem and social support, and decreased depression, substance abuse, and suicidality [55,57,58]. Other surveys found an association between structural stigma at the state level (eg, lack of nondiscrimination policies for sexual minoritized status) and disparities in drug and cigarette use [59,60].

Minority stress models propose that being minoritized is linked to bias, discrimination, and lack of support that can lead to stress, anxiety, and depression, which contribute to adverse long-term psychosocial and health outcomes ([figure 2](#)) [34,61-63]. Structural stigma in the sociocultural environment may interact with individual psychologic characteristics, impacting decision making and behaviors among young sexual minoritized individuals [64]. Whether or not the minority stress model is correct, multiple observational studies demonstrate that rates of mental health problems (eg, anxiety, depression, suicide and self-harm, conduct disorder, substance abuse, and posttraumatic stress disorder) are increased among sexual minoritized youth compared with nonsexual minoritized youth [65-71].

Potential protective factors — Factors that contribute to resilience and counteract stigmatization include [45,64,72-80]:

- Acceptance
- Competence
- Higher levels of self-esteem and psychologic well-being
- Strong sense of self and self-acceptance
- Strong ethnic identification
- Strong connections to family and school
- Caring adult role models outside the family
- Community involvement

Personal, family, and societal acceptance of sexual identity may provide protection against

some of the adverse psychosocial and health outcomes for sexual minoritized youth. Family acceptance is particularly salient [2,3,35]. In multiple observational studies, parental support and acceptance of sexual minoritized youth are correlated with positive mental health outcomes, and parental rejection and lack of support are correlated with negative mental health outcomes [58,81].

Health care providers may contribute to resilience by serving as role models and demonstrating openness, support, and respect for diversity. They also can provide information and a safe place for parent and child discussion and help parents overcome their fears and biases to more fully accept and nurture their sexual minoritized children. (See '[Parents and family](#)' below.)

Health risks — Sexual minoritized youth are more likely than their heterosexual peers to report health risk behaviors, including attempted suicide, substance use, high-risk sexual behaviors, and unhealthy eating or exercise patterns, and experience of victimization/violence [32,36,42,43,69,70,82-84]. Adverse social experience, including discrimination and victimization, have been associated with health problems, sexual risk taking, and HIV exposure [85]. Some risk behaviors appear to vary depending upon subgroups, but the data are limited [44,86-89].

Victimization and violence — Sexual minoritized youth are at higher risk than their nonsexual minoritized peers for sexual prejudice ("homophobia"); verbal harassment; physical threats, harassment, or assault; and physical and sexual abuse [90]. Victimization is an important mediator of psychologic distress and mental health outcomes, including suicidality, among sexual minoritized youth [39,47,91-96]. Sexual minoritized youth may experience victimization and violence at home, school, or in the community [32,37,44,97-99]. Victimization and psychologic distress appear to decrease with age and over time [96,100].

In a 2011 meta-analysis of school-based studies, sexual minoritized individuals were 1.2 times more likely to report parental physical abuse and 3.8 times more likely to report sexual abuse (the site of abuse and perpetrator were not addressed) than their nonsexual minoritized peers [44]; they also were 1.7 times more likely to have been assaulted at school and 2.4 times more likely to miss school out of fear.

In a nationwide survey of high school students, 28 percent reported being electronically bullied; 19 percent reported having been forced to have sexual intercourse at some point

in their lives; and among those who had dated or went out with someone during the 12 months before the survey, 16 percent reported sexual dating violence and 13 percent reported physical dating violence [32]. These rates were generally approximately twice as high as those among heterosexual students.

School victimization — In a 2019 national survey of high school students, sexual minoritized students were more likely than their heterosexual peers to report being threatened or injured with a weapon (12 versus 6 percent) or bullied (32 versus 17 percent) on school property in the previous 12 months or to report not going to school for at least 1 of the previous 30 days because of safety concerns (14 versus 8 percent) [32]

In a 2011 national survey of 8584 lesbian, gay, bisexual, and transgender students (13 to 20 years) [37]:

- 71 to 85 percent reported hearing sexual prejudicial remarks (eg, "that's so gay," "dyke," "faggot") frequently or often at school; 57 percent reported hearing such remarks from teachers or other staff
- 64 percent felt unsafe at school because of their sexual orientation
- 82 percent were verbally harassed (called names or threatened), 38 percent were physically harassed (eg, pushed or shoved), and 18 percent were physically assaulted (eg, punched, kicked, injured with a weapon) at school in the past year because of their sexual orientation
- 55 percent were harassed or threatened by their peers via electronic mediums (eg, text messages, postings on Facebook)
- 32 percent missed at least one day of school in the past month because they felt unsafe or uncomfortable

Middle school sexual minoritized youth were more likely than high school sexual minoritized youth to experience harassment and assault based on sexual orientation or gender expression [37]. School characteristics associated with discrimination included public or religious schools (versus private/nonreligious schools), location in the Southern or Midwestern United States, and location in small town or rural areas.

Rates of bullying and victimization of lesbian, gay, and bisexual youth decreased with

increasing age, as demonstrated in a prospective survey (2004 to 2010) from England [100]. However, there were differences between males and females: By the final wave of the survey, lesbian/bisexual females were no more likely to report bullying than their heterosexual peers, whereas gay/bisexual males were more likely to report bullying than their heterosexual peers.

School victimization is associated with truancy, feeling unsafe, and decreased school engagement and academic achievement, as well as long-term health and behavioral risk factors, emphasizing the importance of early interventions that decrease school victimization and promote life-long educational attainment [101-104]. (See '[School](#)' below.)

Sexual victimization — Rates of sexual violence and victimization (eg, dating violence, child and adolescent sexual abuse) are higher in sexual minoritized individuals than in their peers [105-112]. Adolescents who identify as having partners of both sexes seem to be at highest risk [36]. In a 2011 systematic review of 75 studies (139,635 participants) evaluating sexual assault against sexual minoritized persons, lifetime sexual assault ranged from 16 to 85 percent for women and 12 to 54 percent for men [110]. Lesbian/bisexual women were more likely to report childhood sexual assault, adult sexual assault, and intimate partner violence than gay/bisexual men. Gay/bisexual men were more likely to report hate-crime related assault than lesbian/bisexual women.

Mental health and self-harm — Adolescence is a challenging time for all youth. Sexual minoritized youth have the same mental health needs as their heterosexual peers but face the additional challenge of societal bias against sexual and gender minorities.

Sexual minoritized youth report higher levels of social isolation, low self-esteem, impaired self-concept, and a variety of internalizing (eg, anxiety, depression) and/or externalizing (eg, aggression) symptoms than their peers [41,113-119]. These symptoms may be induced by stigmatization and minority stress or victimization [100]. (See '[Stigmatization and minority stress](#)' above.)

Sexual minoritized youth also report higher levels of suicidality and nonsuicidal self-injury than their peers [32,38,69,120,121]. In a 2018 meta-analysis of 22 observational studies from 10 countries, the risk of suicide attempt was greater in sexual minoritized than heterosexual youth (23.7 versus 6.4 percent, odds ratio [OR] 3.5, 95% CI 3.0-4.1); in subgroup analysis, the risk was greater among bisexual than homosexual youth (OR 4.9 versus 3.7) [87]. In a national survey of high school students in the United States, the

prevalence of nonsuicidal self-injury ranged from 38 to 53 percent among sexual minoritized youth between 2005 and 2017 (compared with 11 to 20 percent among heterosexual youth) [[121](#)].

In the National Longitudinal Study of Adolescent Health, symptoms of depression among sexual minoritized youth decreased between adolescence (1994-1995) and young adulthood (2007-2008), but suicidality did not [[122](#)].

Use of multiple measures of sexual minoritized status and factors related to racial/ethnic background, religious affiliation, and self-identification may help to explain within-group disparities in rates of mental health problems (eg, depression, substance use, eating disorders) and suicidal ideation [[123-125](#)].

Tobacco and substance use — Sexual minoritized youth appear to be at increased risk for tobacco, alcohol, and drug use compared with their nonsexual minoritized peers, but the risk varies with the subgroup, substance, and other factors such as race/ethnicity, socioeconomic status, victimization, and family support [[70,71,126-133](#)].

In a national survey, female sexual minoritized youth (self-categorized as lesbian, bisexual, or "mostly heterosexual") and "mostly heterosexual" males were at least two times more likely to use tobacco than heterosexual youth; tobacco use was not increased among self-categorized gay/bisexual boys [[134](#)]. In other national surveys, bisexual youth report greater substance use than homosexual youth [[70,71,135,136](#)].

Tobacco and/or substance use may be a marker for other risky health behaviors (eg, exchange sex, sex with drug users, multiple partners, unprotected sex) [[137-139](#)]. In a survey of young men who have sex with men (MSM) in New York City, tobacco use was associated with increased use of illicit substances, alcohol, abuse of prescription drugs, and more casual and transactional sex partners [[137](#)]. Young MSM tobacco users were also more likely to use substances before or during sex. In a cross-sectional survey of young MSM, use of methamphetamine was associated with use of other illicit drugs, sex with drug users, multiple partners, lower rates of condom use, and higher rates of sexually transmitted infections [[138](#)].

Homelessness — Sexual minoritized youth who are rejected by their parents and families may run away or be forced to leave home [[140](#)]. In convenience samples of homeless youth, sexual minoritized youth are disproportionately represented [[141,142](#)]. In

two surveys (2005 and 2007) of 6317 high school students from one state, 25 percent of self-identified lesbian/gay and 15 percent of self-identified bisexual youth were homeless, compared with 3 percent of heterosexual youth [\[143\]](#).

Homeless youth may be forced to engage in risky behaviors to survive [\[140\]](#). Compared with nonsexual minoritized peers, runaway and homeless sexual minoritized youth report higher rates of substance use, suicide attempts, risky sexual behaviors (eg, prostitution, survival sex, unprotected intercourse), and sexual victimization [\[144-150\]](#).

Sexually transmitted infections — Although many adolescents become sexually active during high school, sexual minoritized youth are more likely to report engaging in behaviors, or being forced to engage in behaviors, that increase their risk of sexually transmitted infections (eg, multiple partners, lack of condom use, using alcohol or drugs before sexual activity) [\[33,83,151-155\]](#). Sexual minoritized youth may begin experimenting with sex earlier than their heterosexual peers (ie, before age 13 years), use alcohol or drugs with intercourse, lack medically accurate information regarding same-sex safer sex, and lack social support or access to services [\[33,105\]](#). In a meta-analysis of six studies, sexual minoritized youth were almost twice as likely to report sex while intoxicated as heterosexual peers [\[156\]](#).

Young MSM account for a disproportionate number of new HIV infections in the United States [\[157,158\]](#). This is particularly true for those who are Black/African American, Hispanic/Latino, homeless, or engage in survival sex [\[159,160\]](#). New HIV diagnoses attributed to sexual contact among men who have sex with men is highest among young MSM (ie, ages 13 to 24 years) [\[161\]](#). (See ["The adolescent with HIV infection", section on 'Epidemiology'](#).)

Young MSM and young transfeminine youth who engage in receptive anal sex, have multiple partners, or have a partner who is known to be HIV positive may be candidates for HIV pre-exposure prophylaxis (PrEP). Additional considerations, including assessment of the risk of acquiring HIV and the ability to adhere to PrEP, are discussed separately. (See ["Patient evaluation and selection for HIV pre-exposure prophylaxis"](#).)

Young women who have sex with women also may be at risk for HIV depending upon their sexual history and behaviors (eg, coerced sexual contact; exchange sex; digital-vaginal or digital-anal contact, particularly with shared penetrative sex items) [\[7,162-165\]](#).

Unplanned pregnancy — Unplanned pregnancy may occur in sexual minoritized females and transmasculine youth and young adults. Sexual minoritized females may have sex with males as they explore their sexual identity. They also may engage in heterosexual dating and sexual behaviors to avoid being identified as lesbian or bisexual, may engage in exchange sex, or may have coerced sexual contact [7,166-169]. (See "[Contraception: Counseling and selection](#)" and "[Emergency contraception](#)".)

Several population-based surveys have documented higher rates of pregnancy involvement among lesbian, gay, and bisexual youth than their heterosexual peers [82,167,170-173]. In one survey, lesbian and bisexual respondents were as likely to have had penile-vaginal intercourse as heterosexual or unsure adolescents; they were more likely to have used an ineffective method of contraception or no method of contraception and to become pregnant [166]. Adolescents self-identified as lesbians also reported higher rates of a history of sexual abuse and prostitution [174]. In the National Survey of Family Growth (2006 to 2010), lesbian and bisexual young women (15 to 20 years of age) reported earlier heterosexual debut, more male sexual partners, more female sexual partners, and more forced sexual encounters by a male partner than heterosexual young women [82,173].

Weight control and anabolic steroids — Sexual minoritized youth are more likely than their heterosexual peers to report unhealthy weight control behaviors (eg, not eating for ≥ 24 hours; taking diet pills, powders, or liquids; vomiting or taking laxatives) and misuse of anabolic-androgenic steroids [33,175,176].

Patterns of disordered eating appear to differ according to subgroup, but data are limited [177-182]. Male sexual minoritized youth are more likely to report symptoms of eating disorders (dissatisfaction and idealized body image, dieting, bingeing and purging, using weight control products) than heterosexual male peers [179-181]. Female sexual minoritized youth are more likely to report bingeing with or without purging than their heterosexual female peers.

Male sexual minoritized youth also are more likely to misuse anabolic-androgenic steroids than their heterosexual peers. In a national survey of high school students (2005 to 2007), misuse of anabolic-androgenic steroids was more common among sexual minoritized than heterosexual boys (21 versus 4 percent, OR 5.8, 95% CI 4.1-8.2) [175]. Secondary analysis suggested that depression/suicidality, victimization, and substance use

contributed to misuse of steroids; body dissatisfaction was not assessed.

Unintentional injury — Other risk behaviors that have been more frequently reported among sexual minoritized than nonsexual minoritized youth in the Youth Risk Behavior Surveillance System include behaviors that contribute to unintentional injury (eg, rarely/never wore seatbelt, rode with a driver who had been drinking alcohol, drove when drinking alcohol) [[33,183](#)].

SUPPORT AND ADVOCACY

Medical providers are in a unique position to provide and model positive regard, support, and open acceptance for youth of all sexual orientations and identities. Health care providers who accept and support gender or sexual diversity can serve as models for parents and other family members. Negative messaging from adults about sexual minorities may be subtle or overt, but is picked up by youth of all ages. Prepubertal children are particularly sensitive to social cues and often learn to hide their interests, preferences, and play after getting negative feedback from peers and adults.

Normalization of sexuality — Health care professionals who provide developmental screening and anticipatory guidance can integrate conversations around gender and sexuality into their education and counseling. Ongoing conversations about gender and sexuality may allow earlier identification and support for sexual minoritized youth and their families.

- Including questions about gender play and preferences, body image and esteem, expression of femininity and masculinity in routine health care visits from the toddler ([table 1](#) and [table 2](#)) through teenage years ([table 3](#)) normalizes the youth's ongoing and evolving sexuality, a universal developmental experience.

Asking such questions provides a developmentally appropriate introduction to upcoming discussions about puberty and adolescence and can be helpful for both the parents and the child to practice open communication about these sensitive issues within the medical setting. It also allows parents, children, and adolescents, including those who are sexual minoritized or have experienced sexual prejudice, to realize that the medical visit can be a safe place to explore questions and concerns about sexual health.

- The health care provider's asking parents about their child's activities and preferences in the prepubertal years demonstrates that adults should be interested in their child's development of identity and provides appropriate models for communication and support about these issues ([table 1](#)).
- The provider's open and curious approach to understanding the ways children embrace their gender and sexual selves can guide parents in approaching their gender-diverse child or sexual minoritized adolescent.

Early identification of gender or sexual diversity — Early identification of gender or sexual diversity may permit earlier supportive interventions for the child and family. Asking the child and parent(s) about how the child might view their gender, their body, and their femininity or masculinity is usually well tolerated ([table 1](#) and [table 2](#)). Parents may be reassured to hear that most children explore and experiment with gender and sexuality to some extent during childhood. The provider can then encourage the parents to focus on supporting their child as he or she explores and develops his or her sexuality rather than trying to predict future sexual orientation. (See "[Gender development and clinical presentation of gender diversity in children and adolescents](#)", [section on 'Gender development in childhood'](#).)

Obtaining a more detailed exploration of gender and sexuality may be warranted for children and adolescents who present with suicidality, mood, behavior, or school problems. Sexual minoritized children may initially present with depression, anxiety, behavior problems, or school problems rather than explicitly identifying or verbalizing more complex constructs of same-sex sexual attractions or brain and body gender incongruence.

Parents and family — The parents and family of sexual minoritized youth play an important role in healthy adolescent development [[12,57,81,184-186](#)]. However, they may have difficulty accepting having a sexual minoritized child or family member. Their difficulty may be related to fear of the unknown, fear of social stigma, and fear for their child's or relative's safety. These concerns may reflect their own biases and personal, cultural, religious, and historical background. Parents should understand that their concerns may be communicated to the child, both explicitly and implicitly, and affect how the child views himself or herself.

Health care providers can help parents work through their values and beliefs in a way that

consistently supports the child. Education, parent-to-parent support groups, or therapy may be helpful in this regard. In the experience of the authors of this topic review, parents appreciate and value a health care provider who acknowledges their confusion or distress. Providers can guide parents towards resources that may be helpful to them and other family members and help them to focus on creating an accepting and safe environment for their child ([table 4](#)).

It may be helpful to explain that same sex attractions are a normal variant of sexuality; being attracted to members of the same sex is not a mental disorder [[2,3,34,35,187](#)]. Familial support and acceptance of the sexual minoritized youth (ie, affirmation) helps adolescents explore their sexual identities in a safe environment [[35](#)]. Interventions that attempt to change sexual orientation (ie, "reparative" or "conversion" therapy) are ineffective, coercive, and potentially harmful (by increasing internalized stigma, distress, and depression) [[2,34,35,187,188](#)].

Health care providers of sexual minoritized youth can model consistent, positive, and strength-based acceptance and support for parents and family members who want to be more supportive of their sexual minoritized youth. They can also explore and address parental concerns and provide early referrals to community and advocacy resources. (See '[Resources](#)' below.)

Disclosure — Children and adolescents who are sexual minorities may or may not feel initial confusion, discomfort, or emotional turmoil as they come to terms with their sexual attractions, desires, and behaviors.

It is important to allow children, youth, and adults the freedom and autonomy to choose, consciously or unconsciously, the right time to disclose their sexual minoritized status ("come out") to their provider, family members, or peers. The process of coming out can be lifelong, as youth meet new people and experience new social situations. Sexual minoritized youth typically first disclose to a friend [[189](#)]. Most youth tell at least one parent, more often the mother, or they tell the mother before the father (in heterosexual parent dynamics) [[190](#)]. When a sexual minoritized youth has disclosed to a provider, maintaining privacy, confidentiality, and respecting the youth's plan for disclosure is paramount [[2](#)].

Health care providers may be asked to address questions or concerns about disclosure to family or friends [[12](#)]. Discussions about disclosure should include the timing, approach,

and potential repercussions ([table 5](#)). Providers can help adolescents think through the pros and cons of disclosure, whom to tell, how to make the disclosure, and provide other guidance in revealing, asserting, and feeling good about their sexual orientation or other sexual minoritized status.

Given the potential for discrimination and bias against sexual minorities, disclosure can be an enormous challenge. Reactions to disclosure (positive or negative) are not always predictable. Positive and accepting, or negative and rejecting, family reactions have long-term implications on the health and well-being of sexual minoritized youth [\[57,81,184\]](#). (See '[Potential protective factors](#)' above.)

Coming out can be an opportunity as well as a challenge. Potential benefits of disclosure include improved communication, support, and intimacy with family and friends; decreased fears and worry about inadvertent disclosure; increased opportunities to access care, social networks, and resources; more authentic internal and external presentation; and acceptance of one's true self [\[191\]](#). Negative reactions should be discussed and anticipated. Disclosure may increase the rate of victimization [\[191\]](#). The safety of the youth is paramount. There are particular situations where it may not be safe, helpful, or advisable to disclose. Some youth have significant and real concerns regarding physical safety or abuse from a parent or guardian due to sexual minoritized status. Others may not be physically unsafe but may be rejected, emotionally and verbally abused, or coerced financially. Dependence on parents for housing, schooling, food, and other expenses is a real concern for many youth. If a youth is uncertain how a parent or guardian will react, it may be safer to wait to disclose until the youth is in a more independent or supported position. Disproportionate numbers of homeless youth are sexual minorities, suggesting that many were rejected or not safe at home in their asserted sexual identity [\[144,192,193\]](#). (See '[Homelessness](#)' above.)

Health care providers can help sexual minoritized youth who do not have supportive families identify and engage in other social and support networks (eg, gay-straight alliances [GSAs], community organizations). It is helpful for providers to be familiar with local groups or agencies that offer a positive environment and support for sexual minoritized youth; providers may identify local groups through national organizations or networking with local providers or groups ([table 6](#)). It is also helpful for the agencies to know which providers in the community provide safe and appropriate health services for sexual minoritized youth. (See '[Resources](#)' below and '[Lesbian, gay, bisexual, and other](#)

[sexual minoritized youth: Primary care", section on 'Resources'.\)](#)

School — Difficult school environments for sexual minoritized youth may result in lower educational achievement and increased risk for depression and suicide [37,48,194]. Schools that institute specific policies and programs promoting safety and diversity create an opportunity for all students to achieve both academic and social goals [195].

A tolerant and supportive school climate includes a leadership plan with staff who are trained, committed, and can implement strategies designed to promote zero tolerance of harassment and reduce victimization. School personnel who promote tolerance of diversity improve sexual minoritized students' feelings of connectedness and safety [196]. Protective school environments have been correlated with lower risks of substance use and suicidality among sexual minoritized youth [48,197].

Schools can promote a safer climate for sexual minoritized youth when [100,170,198-201]:

- Antidiscrimination and antiharassment policies include sexual minoritized youth and are strongly enforced, with staff actively intervening to enforce the policies whenever it is necessary.
- They have identified an individual (eg, guidance counselor, principal) and/or a process to address concerns of sexual minoritized youth.
- There are social opportunities such as GSAs integrated into the school setting; GSAs are student-led clubs that provide a place for sexual minoritized youth to gather, socialize, and educate in a safe and supportive context. The [GSAnetwork](#) can provide resources for students who would like to start a GSA. Participation in and perceived effectiveness of a GSA may decrease victimization and its effects on well-being [202].
- Sexual minoritized issues are integrated into the larger school curriculums (eg, discussion of same-sex relationships in sex education courses, discussion of sexual prejudice in athletics programs and physical education classes) [170,203].

A detailed discussion of policy recommendations to increase the safety of sexual minoritized youth at school is beyond the scope of this review but is available in [Safe at School: Addressing the School Environment and LGBT safety through Policy and Legislation](#) [194].

Resources — The tables provide lists of resources that may be helpful to sexual minoritized youth ([table 6](#)), parents and family members ([table 4](#)), and clinicians ([table 7](#)).

SOCIETY GUIDELINE LINKS

Links to society and government-sponsored guidelines from selected countries and regions around the world are provided separately. (See "[Society guideline links: Adolescent sexual health and pregnancy](#)" and "[Society guideline links: Sexually transmitted infections](#)" and "[Society guideline links: HIV infection in adolescents](#)" and "[Society guideline links: Contraception](#)" and "[Society guideline links: Medical care for homeless persons](#)".)

SUMMARY

- Sexual orientation has multiple dimensions, including attractions, identity, behaviors, and the gender(s) of partners. Sexual identity is an individual's assessment of his or her sexual orientation. It is best defined by the individual adolescent; many avoid discreet labels. (See '[Terminology](#)' above.)
- The term "sexual minoritized" encompasses a variety of gender and sexual identities and expressions that differ from cultural norms. In this topic, we focus on youth who identify themselves as lesbian, gay, bisexual, pansexual, or unsure of their sexual identity, who avoid discrete sexual orientation labels. (See '[Terminology](#)' above.)
- Adolescents explore their emerging sexuality through attractions, fantasies, and behaviors. Adolescent sexual identity and self-identified orientation does not necessarily correlate with sexual behaviors. (See '[Developmental perspective](#)' above.)
- In large surveys, 4 to 7 percent of high school youth report same-sex attractions or behaviors. (See '[Epidemiology](#)' above.)
- The majority of sexual minoritized youth are healthy and well adjusted. However, compared with heterosexual peers, sexual minoritized youth are at increased risk of adverse psychosocial and health outcomes, including victimization, depression, suicide, substance use, homelessness, sexually transmitted infections, and unplanned

pregnancy. (See ['Potential psychosocial and health concerns'](#) above.)

- Adverse psychosocial and health outcomes may be related to stigmatization and lack of acceptance by individuals, communities, and cultures ([figure 2](#)). Factors that may counteract stigmatization include acceptance, competence, increased self-esteem, strong connections to family and school, caring adult role models, and involvement in a community. (See ['Stigmatization and minority stress'](#) above and ['Potential protective factors'](#) above.)
- Health care providers can support sexual minoritized youth by normalizing sexuality throughout childhood; early identification of gender or sexual diversity and associated psychosocial and health concerns ([table 1](#) and [table 2](#)); helping parents support their child; helping the youth with decisions about disclosure ([table 5](#)); advocating for schools that provide safety for sexual minoritized youth; and providing educational and support resources to the sexual minoritized youth ([table 6](#)) and his or her family ([table 4](#)). (See ['Support and advocacy'](#) above.)

Use of UpToDate is subject to the [Subscription and License Agreement](#).

REFERENCES

1. [Goldfried MR, Bell AC. Extending the boundaries of research on adolescent development. J Clin Child Adolesc Psychol 2003; 32:531.](#)
2. [Committee On Adolescence. Office-based care for lesbian, gay, bisexual, transgender, and questioning youth. Pediatrics 2013; 132:198.](#)
3. [Adelson SL, American Academy of Child and Adolescent Psychiatry \(AACAP\) Committee on Quality Issues \(CQI\). Practice parameter on gay, lesbian, or bisexual sexual orientation, gender nonconformity, and gender discordance in children and adolescents. J Am Acad Child Adolesc Psychiatry 2012; 51:957.](#)
4. [Spigarelli MG. Adolescent sexual orientation. Adolesc Med State Art Rev 2007; 18:508.](#)
5. [Austin SB, Conron K, Patel A, Freedner N. Making sense of sexual orientation measures: findings from a cognitive processing study with adolescents on health survey questions. J LGBT Health Res 2007; 3:55.](#)
6. [Chandra A, Mosher WD, Copen C, Sionean C. Sexual behavior, sexual attraction, and sexual identity in the United States: data from the 2006-2008 National Survey of](#)

[Family Growth. Natl Health Stat Report 2011; :1.](#)

7. [Goodenow C, Szalacha LA, Robin LE, Westheimer K. Dimensions of sexual orientation and HIV-related risk among adolescent females: evidence from a statewide survey. Am J Public Health 2008; 98:1051.](#)
8. [Mustanski B, Birkett M, Greene GJ, et al. The association between sexual orientation identity and behavior across race/ethnicity, sex, and age in a probability sample of high school students. Am J Public Health 2014; 104:237.](#)
9. [Igartua K, Thombs BD, Burgos G, Montoro R. Concordance and discrepancy in sexual identity, attraction, and behavior among adolescents. J Adolesc Health 2009; 45:602.](#)
10. [McCabe J, Brewster KL, Tillman KH. Patterns and correlates of same-sex sexual activity among U.S. teenagers and young adults. Perspect Sex Reprod Health 2011; 43:142.](#)
11. [Pathela P, Hajat A, Schillinger J, et al. Discordance between sexual behavior and self-reported sexual identity: a population-based survey of New York City men. Ann Intern Med 2006; 145:416.](#)
12. [Bell DL, Breland DJ, Ott MA. Adolescent and young adult male health: a review. Pediatrics 2013; 132:535.](#)
13. [Coker TR, Austin SB, Schuster MA. Health and healthcare for lesbian, gay, bisexual, and transgender youth: reducing disparities through research, education, and practice. J Adolesc Health 2009; 45:213.](#)
14. [Coker TR, Austin SB, Schuster MA. The health and health care of lesbian, gay, and bisexual adolescents. Annu Rev Public Health 2010; 31:457.](#)
15. [Bullough VL. Children and adolescents as sexual beings: a historical overview. Child Adolesc Psychiatr Clin N Am 2004; 13:447.](#)
16. Kinsey AC, Pomeroy WB, Martin CE. Sexual behavior in the human male, WB Saunders , Philadelphia 1948.
17. [Levine DA, Committee On Adolescence. Office-based care for lesbian, gay, bisexual, transgender, and questioning youth. Pediatrics 2013; 132:e297.](#)
18. Green R. The "Sissy Boy Syndrome" and the Development of Homosexuality, Yale University Press, New Haven, CT 1987.
19. [Money J, Russo AJ. Homosexual outcome of discordant gender identity/role in childhood: Longitudinal follow-up. J Pediatr Psychol 1979; 4:29.](#)
20. Zucker KJ, Bradley SJ. Gender Identity Disorder and Psychosexual Problems in Childre

n and Adolescents, Guilford Press, New York 1995.

21. [Zuger B. Early effeminate behavior in boys. Outcome and significance for homosexuality. J Nerv Ment Dis 1984; 172:90.](#)
22. [Ryan C, Futterman D. Lesbian and Gay Youth: Care and Counseling. Adolesc Med 1997; 8:207.](#)
23. [Remafedi G, Resnick M, Blum R, Harris L. Demography of sexual orientation in adolescents. Pediatrics 1992; 89:714.](#)
24. [Berona J, Stepp SD, Hipwell AE, Keenan KE. Trajectories of Sexual Orientation from Adolescence to Young Adulthood: Results from a Community-Based Urban Sample of Girls. J Adolesc Health 2018; 63:57.](#)
25. [Rosario M, Schrimshaw EW, Hunter J. Predicting different patterns of sexual identity development over time among lesbian, gay, and bisexual youths: a cluster analytic approach. Am J Community Psychol 2008; 42:266.](#)
26. [Fasula AM, Oraka E, Jeffries WL 4th, et al. Young Sexual Minority Males in the United States: Sociodemographic Characteristics And Sexual Attraction, Identity and Behavior. Perspect Sex Reprod Health 2016; 48:3.](#)
27. [Mallon GP, DeCrescenzo T. Transgender children and youth: a child welfare practice perspective. Child Welfare 2006; 85:215.](#)
28. [Plummer DC. The quest for modern manhood: masculine stereotypes, peer culture and the social significance of homophobia. J Adolesc 2001; 24:15.](#)
29. [Piper J, Mannino M. Identity formation for individuals in transition: A narrative family therapy model. J GLBT Fam Stud 2008; 4:75.](#)
30. [Moser C, Kleinplatz PJ, Zuccarini D, Reiner WG. Situating unusual child and adolescent sexual behavior in context. Child Adolesc Psychiatr Clin N Am 2004; 13:569.](#)
31. [Parmar DD, Alabaster A, Vance S Jr, et al. Identification of Sexual Minority Youth in Pediatric Primary Care Settings Within a Large Integrated Healthcare System Using Electronic Health Records. J Adolesc Health 2020; 66:255.](#)
32. [Johns MM, Lowry R, Haderxhanaj LT, et al. Trends in Violence Victimization and Suicide Risk by Sexual Identity Among High School Students - Youth Risk Behavior Survey, United States, 2015-2019. MMWR Suppl 2020; 69:19.](#)
33. Centers for Disease Control and Prevention. High School YRBS. Youth Online. Supplementary tables. https://www.cdc.gov/healthyyouth/data/yrbs/2019_tables/index.htm (

Accessed on September 09, 2020).

34. IOM (Institute of Medicine). The Health of Lesbian, Gay, Bisexual, and Transgender People: Building a Foundation for Better Understanding, National Academies Press, Washington, DC 2011.
35. [Society for Adolescent Health and Medicine. Recommendations for promoting the health and well-being of lesbian, gay, bisexual, and transgender adolescents: a position paper of the Society for Adolescent Health and Medicine. J Adolesc Health 2013; 52:506.](#)
36. [Pathela P, Schillinger JA. Sexual behaviors and sexual violence: adolescents with opposite-, same-, or both-sex partners. Pediatrics 2010; 126:879.](#)
37. Kosciw JG, Greytak EA, Bartkiewicz MJ, et al. The 2011 National School Climate Survey. The experiences of lesbian, gay, bisexual, and transgender youth in our nation's schools. <http://www.glsen.org/sites/default/files/2011%20National%20School%20Climate%20Survey%20Full%20Report.pdf> (Accessed on November 02, 2013).
38. [Marshal MP, Dietz LJ, Friedman MS, et al. Suicidality and depression disparities between sexual minority and heterosexual youth: a meta-analytic review. J Adolesc Health 2011; 49:115.](#)
39. [Oshri A, Handley ED, Sutton TE, et al. Developmental trajectories of substance use among sexual minority girls: associations with sexual victimization and sexual health risk. J Adolesc Health 2014; 55:100.](#)
40. [Strutz KL, Herring AH, Halpern CT. Health disparities among young adult sexual minorities in the U.S. Am J Prev Med 2015; 48:76.](#)
41. [Luk JW, Gilman SE, Haynie DL, Simons-Morton BG. Sexual Orientation and Depressive Symptoms in Adolescents. Pediatrics 2018; 141.](#)
42. [Ivey-Stephenson AZ, Demissie Z, Crosby AE, et al. Suicidal Ideation and Behaviors Among High School Students - Youth Risk Behavior Survey, United States, 2019. MMWR Suppl 2020; 69:47.](#)
43. [Jones CM, Clayton HB, Deputy NP, et al. Prescription Opioid Misuse and Use of Alcohol and Other Substances Among High School Students - Youth Risk Behavior Survey, United States, 2019. MMWR Suppl 2020; 69:38.](#)
44. [Friedman MS, Marshal MP, Guadamuz TE, et al. A meta-analysis of disparities in childhood sexual abuse, parental physical abuse, and peer victimization among](#)

- [sexual minority and sexual nonminority individuals. Am J Public Health 2011; 101:1481.](#)
45. [Herrick AL, Egan JE, Coulter RW, et al. Raising sexual minority youths' health levels by incorporating resiliencies into health promotion efforts. Am J Public Health 2014; 104:206.](#)
 46. [Herrick AL, Lim SH, Plankey MW, et al. Adversity and syndemic production among men participating in the multicenter AIDS cohort study: a life-course approach. Am J Public Health 2013; 103:79.](#)
 47. [Duncan DT, Hatzenbuehler ML. Lesbian, gay, bisexual, and transgender hate crimes and suicidality among a population-based sample of sexual-minority adolescents in Boston. Am J Public Health 2014; 104:272.](#)
 48. [Hatzenbuehler ML, Birkett M, Van Wagenen A, Meyer IH. Protective school climates and reduced risk for suicide ideation in sexual minority youths. Am J Public Health 2014; 104:279.](#)
 49. [Eccles TA, Sayegh MA, Fortenberry JD, Zimet GD. More normal than not: a qualitative assessment of the developmental experiences of gay male youth. J Adolesc Health 2004; 35:425.e11.](#)
 50. [Friedman MS, Marshal MP, Stall R, et al. Gay-related development, early abuse and adult health outcomes among gay males. AIDS Behav 2008; 12:891.](#)
 51. [Frost DM, Parsons JT, Nanín JE. Stigma, concealment and symptoms of depression as explanations for sexually transmitted infections among gay men. J Health Psychol 2007; 12:636.](#)
 52. [Alderman EM, Breuner CC, COMMITTEE ON ADOLESCENCE. Unique Needs of the Adolescent. Pediatrics 2019; 144.](#)
 53. [Bruce D, Stall R, Fata A, Campbell RT. Modeling minority stress effects on homelessness and health disparities among young men who have sex with men. J Urban Health 2014; 91:568.](#)
 54. [Austin A, Herrick H, Proescholdbell S. Adverse Childhood Experiences Related to Poor Adult Health Among Lesbian, Gay, and Bisexual Individuals. Am J Public Health 2016; 106:314.](#)
 55. [Veale JF, Peter T, Travers R, Saewyc EM. Enacted Stigma, Mental Health, and Protective Factors Among Transgender Youth in Canada. Transgend Health 2017; 2:207.](#)

56. [Clark KA, Cochran SD, Maiolatesi AJ, Pachankis JE. Prevalence of Bullying Among Youth Classified as LGBTQ Who Died by Suicide as Reported in the National Violent Death Reporting System, 2003-2017. JAMA Pediatr 2020; 174:1211.](#)
57. [Ryan C, Huebner D, Diaz RM, Sanchez J. Family rejection as a predictor of negative health outcomes in white and Latino lesbian, gay, and bisexual young adults. Pediatrics 2009; 123:346.](#)
58. [Ryan C, Russell ST, Huebner D, et al. Family acceptance in adolescence and the health of LGBT young adults. J Child Adolesc Psychiatr Nurs 2010; 23:205.](#)
59. [Hatzenbuehler ML, Jun HJ, Corliss HL, Bryn Austin S. Structural stigma and sexual orientation disparities in adolescent drug use. Addict Behav 2015; 46:14.](#)
60. [Hatzenbuehler ML, Jun HJ, Corliss HL, Austin SB. Structural stigma and cigarette smoking in a prospective cohort study of sexual minority and heterosexual youth. Ann Behav Med 2014; 47:48.](#)
61. [Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. Psychol Bull 2003; 129:674.](#)
62. [Frost DM, Lehavot K, Meyer IH. Minority stress and physical health among sexual minority individuals. J Behav Med 2015; 38:1.](#)
63. [Shilo G, Mor Z. The impact of minority stressors on the mental and physical health of lesbian, gay, and bisexual youths and young adults. Health Soc Work 2014; 39:161.](#)
64. [Pachankis JE, Hatzenbuehler ML, Starks TJ. The influence of structural stigma and rejection sensitivity on young sexual minority men's daily tobacco and alcohol use. Soc Sci Med 2014; 103:67.](#)
65. [Eliason M. Introduction to special issue on suicide, mental health, and youth development. J Homosex 2011; 58:4.](#)
66. [Haas AP, Eliason M, Mays VM, et al. Suicide and suicide risk in lesbian, gay, bisexual, and transgender populations: review and recommendations. J Homosex 2011; 58:10.](#)
67. [Mustanski BS, Garofalo R, Emerson EM. Mental health disorders, psychological distress, and suicidality in a diverse sample of lesbian, gay, bisexual, and transgender youths. Am J Public Health 2010; 100:2426.](#)
68. [King M, Semlyen J, Tai SS, et al. A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people. BMC Psychiatry 2008; 8:70.](#)
69. [Stone DM, Luo F, Ouyang L, et al. Sexual orientation and suicide ideation, plans,](#)

- [attempts, and medically serious attempts: evidence from local Youth Risk Behavior Surveys, 2001-2009. Am J Public Health 2014; 104:262.](#)
70. [Newcomb ME, Birkett M, Corliss HL, Mustanski B. Sexual orientation, gender, and racial differences in illicit drug use in a sample of US high school students. Am J Public Health 2014; 104:304.](#)
71. [Fish JN, Baams L. Trends in Alcohol-Related Disparities Between Heterosexual and Sexual Minority Youth from 2007 to 2015: Findings from the Youth Risk Behavior Survey. LGBT Health 2018; 5:359.](#)
72. [Bird JD, Kuhns L, Garofalo R. The impact of role models on health outcomes for lesbian, gay, bisexual, and transgender youth. J Adolesc Health 2012; 50:353.](#)
73. [Eisenberg ME, Resnick MD. Suicidality among gay, lesbian and bisexual youth: the role of protective factors. J Adolesc Health 2006; 39:662.](#)
74. [Cohn TJ, Hastings SL. Resilience among rural lesbian youth. J Lesbian Stud 2010; 14:71.](#)
75. [Russell ST, Muraco A, Subramaniam A, Laub C. Youth empowerment and high school Gay-Straight Alliances. J Youth Adolesc 2009; 38:891.](#)
76. [Turanovic JJ, Pratt TC. Longitudinal effects of violent victimization during adolescence on adverse outcomes in adulthood: a focus on prosocial attachments. J Pediatr 2015; 166:1062.](#)
77. [Hussen SA, Harper GW, Bauermeister JA, et al. Psychosocial influences on engagement in care among HIV-positive young black gay/bisexual and other men who have sex with men. AIDS Patient Care STDS 2015; 29:77.](#)
78. [Whitaker K, Shapiro VB, Shields JP. School-Based Protective Factors Related to Suicide for Lesbian, Gay, and Bisexual Adolescents. J Adolesc Health 2016; 58:63.](#)
79. [Steiner RJ, Sheremenko G, Lesesne C, et al. Adolescent Connectedness and Adult Health Outcomes. Pediatrics 2019; 144.](#)
80. [Perales F, Campbell A. Health Disparities Between Sexual Minority and Different-Sex-Attracted Adolescents: Quantifying the Intervening Role of Social Support and School Belonging. LGBT Health 2020; 7:146.](#)
81. [Bouris A, Guilamo-Ramos V, Pickard A, et al. A systematic review of parental influences on the health and well-being of lesbian, gay, and bisexual youth: time for a new public health research and practice agenda. J Prim Prev 2010; 31:273.](#)

82. [Tornello SL, Riskind RG, Patterson CJ. Sexual orientation and sexual and reproductive health among adolescent young women in the United States. J Adolesc Health 2014; 54:160.](#)
83. [Thurston IB, Dietrich J, Bogart LM, et al. Correlates of sexual risk among sexual minority and heterosexual South African youths. Am J Public Health 2014; 104:1265.](#)
84. [Amos R, Manalastas EJ, White R, et al. Mental health, social adversity, and health-related outcomes in sexual minority adolescents: a contemporary national cohort study. Lancet Child Adolesc Health 2020; 4:36.](#)
85. [Herrick A, Stall R, Egan J, et al. Pathways towards risk: syndemic conditions mediate the effect of adversity on HIV risk behaviors among young men who have sex with men \(YMSM\). J Urban Health 2014; 91:969.](#)
86. [Lyons A, Hosking W. Health disparities among common subcultural identities of young gay men: physical, mental, and sexual health. Arch Sex Behav 2014; 43:1621.](#)
87. [di Giacomo E, Krausz M, Colmegna F, et al. Estimating the Risk of Attempted Suicide Among Sexual Minority Youths: A Systematic Review and Meta-analysis. JAMA Pediatr 2018; 172:1145.](#)
88. [Rasberry CN, Lowry R, Johns M, et al. Sexual Risk Behavior Differences Among Sexual Minority High School Students - United States, 2015 and 2017. MMWR Morb Mortal Wkly Rep 2018; 67:1007.](#)
89. [Johns MM, Lowry R, Rasberry CN, et al. Violence Victimization, Substance Use, and Suicide Risk Among Sexual Minority High School Students - United States, 2015-2017. MMWR Morb Mortal Wkly Rep 2018; 67:1211.](#)
90. [Baams L. Disparities for LGBTQ and Gender Nonconforming Adolescents. Pediatrics 2018; 141.](#)
91. [Busseri MA, Willoughby T, Chalmers H, Bogaert AF. On the association between sexual attraction and adolescent risk behavior involvement: Examining mediation and moderation. Dev Psychol 2008; 44:69.](#)
92. [Shields JP, Whitaker K, Glassman J, et al. Impact of victimization on risk of suicide among lesbian, gay, and bisexual high school students in San Francisco. J Adolesc Health 2012; 50:418.](#)
93. [Priebe G, Svedin CG. Online or off-line victimisation and psychological well-being: a comparison of sexual-minority and heterosexual youth. Eur Child Adolesc Psychiatry](#)

[2012; 21:569.](#)

94. [van Bergen DD, Bos HM, van Lisdonk J, et al. Victimization and suicidality among Dutch lesbian, gay, and bisexual youths. Am J Public Health 2013; 103:70.](#)
95. [Patrick DL, Bell JF, Huang JY, et al. Bullying and quality of life in youths perceived as gay, lesbian, or bisexual in Washington State, 2010. Am J Public Health 2013; 103:1255.](#)
96. [Birkett M, Newcomb ME, Mustanski B. Does it get better? A longitudinal analysis of psychological distress and victimization in lesbian, gay, bisexual, transgender, and questioning youth. J Adolesc Health 2015; 56:280.](#)
97. [O'Malley Olsen E, Kann L, Vivolo-Kantor A, et al. School violence and bullying among sexual minority high school students, 2009-2011. J Adolesc Health 2014; 55:432.](#)
98. [Schuster MA, Bogart LM, Elliott MN, et al. A longitudinal study of bullying of sexual-minority youth. N Engl J Med 2015; 372:1872.](#)
99. [Caputi TL, Shover CL, Watson RJ. Physical and Sexual Violence Among Gay, Lesbian, Bisexual, and Questioning Adolescents. JAMA Pediatr 2020; 174:791.](#)
100. [Robinson JP, Espelage DL, Rivers I. Developmental trends in peer victimization and emotional distress in LGB and heterosexual youth. Pediatrics 2013; 131:423.](#)
101. [Birkett M, Russell ST, Corliss HL. Sexual-orientation disparities in school: the mediational role of indicators of victimization in achievement and truancy because of feeling unsafe. Am J Public Health 2014; 104:1124.](#)
102. [Rosario M, Corliss HL, Everett BG, et al. Mediation by peer violence victimization of sexual orientation disparities in cancer-related tobacco, alcohol, and sexual risk behaviors: pooled youth risk behavior surveys. Am J Public Health 2014; 104:1113.](#)
103. [Ueno K, Roach TA, Peña-Talamantes AE. The dynamic association between same-sex contact and educational attainment. Adv Life Course Res 2013; 18:127.](#)
104. [Bouris A, Everett BG, Heath RD, et al. Effects of Victimization and Violence on Suicidal Ideation and Behaviors Among Sexual Minority and Heterosexual Adolescents. LGBT Health 2016; 3:153.](#)
105. [Garofalo R, Wolf RC, Kessel S, et al. The association between health risk behaviors and sexual orientation among a school-based sample of adolescents. Pediatrics 1998; 101:895.](#)
106. [Austin SB, Roberts AL, Corliss HL, Molnar BE. Sexual violence victimization history and](#)

- [sexual risk indicators in a community-based urban cohort of "mostly heterosexual" and heterosexual young women. Am J Public Health 2008; 98:1015.](#)
107. [Corliss HL, Cochran SD, Mays VM. Reports of parental maltreatment during childhood in a United States population-based survey of homosexual, bisexual, and heterosexual adults. Child Abuse Negl 2002; 26:1165.](#)
 108. [Saewyc EM, Skay CL, Pettingell SL, et al. Hazards of stigma: the sexual and physical abuse of gay, lesbian, and bisexual adolescents in the United States and Canada. Child Welfare 2006; 85:195.](#)
 109. [D'Augelli AR, Grossman AH, Starks MT. Childhood gender atypicality, victimization, and PTSD among lesbian, gay, and bisexual youth. J Interpers Violence 2006; 21:1462.](#)
 110. [Rothman EF, Exner D, Baughman AL. The prevalence of sexual assault against people who identify as gay, lesbian, or bisexual in the United States: a systematic review. Trauma Violence Abuse 2011; 12:55.](#)
 111. [Cramer RJ, McNiel DE, Holley SR, et al. Mental health in violent crime victims: Does sexual orientation matter? Law Hum Behav 2012; 36:87.](#)
 112. [Walls NE, Atteberry-Ash B, Kattari SK, et al. Gender Identity, Sexual Orientation, Mental Health, and Bullying as Predictors of Partner Violence in a Representative Sample of Youth. J Adolesc Health 2019; 64:86.](#)
 113. [Egan SK, Perry DG. Gender identity: a multidimensional analysis with implications for psychosocial adjustment. Dev Psychol 2001; 37:451.](#)
 114. [Yunger JL, Carver PR, Perry DG. Does gender identity influence children's psychological well-being? Dev Psychol 2004; 40:572.](#)
 115. [Carver PR, Egan SK, Perry DG. Children who question their heterosexuality. Dev Psychol 2004; 40:43.](#)
 116. [Shearer A, Herres J, Kodish T, et al. Differences in Mental Health Symptoms Across Lesbian, Gay, Bisexual, and Questioning Youth in Primary Care Settings. J Adolesc Health 2016; 59:38.](#)
 117. [Becerra-Culqui TA, Liu Y, Nash R, et al. Mental Health of Transgender and Gender Nonconforming Youth Compared With Their Peers. Pediatrics 2018; 141.](#)
 118. [Irish M, Solmi F, Mars B, et al. Depression and self-harm from adolescence to young adulthood in sexual minorities compared with heterosexuals in the UK: a population-based cohort study. Lancet Child Adolesc Health 2019; 3:91.](#)

119. [Bettis AH, Liu RT. Population-Based Analysis of Temporal Trends in the Prevalence of Depressed Mood Among Sexual Minority and Heterosexual Youths From 1999 Through 2017. JAMA Pediatr 2019; 173:1201.](#)
120. [Caputi TL, Smith D, Ayers JW. Suicide Risk Behaviors Among Sexual Minority Adolescents in the United States, 2015. JAMA 2017; 318:2349.](#)
121. [Liu RT. Temporal Trends in the Prevalence of Nonsuicidal Self-injury Among Sexual Minority and Heterosexual Youth From 2005 Through 2017. JAMA Pediatr 2019; 173:790.](#)
122. [Cardom R, Rostosky S, Danner F. Does "it get better" for depressed sexual minority youth in young adulthood? J Adolesc Health 2013; 53:671.](#)
123. [Almazan EP, Roettger ME, Acosta PS. Measures of sexual minority status and suicide risk among young adults in the United States. Arch Suicide Res 2014; 18:274.](#)
124. [Burns MN, Ryan DT, Garofalo R, et al. Mental health disorders in young urban sexual minority men. J Adolesc Health 2015; 56:52.](#)
125. [Lytle MC, De Luca SM, Blosnich JR, Brownson C. Associations of racial/ethnic identities and religious affiliation with suicidal ideation among lesbian, gay, bisexual, and questioning individuals. J Affect Disord 2015; 178:39.](#)
126. [Corliss HL, Rosario M, Birkett MA, et al. Sexual orientation disparities in adolescent cigarette smoking: intersections with race/ethnicity, gender, and age. Am J Public Health 2014; 104:1137.](#)
127. [Kerr D, Ding K, Burke A, Ott-Walter K. An alcohol, tobacco, and other drug use comparison of lesbian, bisexual, and heterosexual undergraduate women. Subst Use Misuse 2015; 50:340.](#)
128. [Kecojevic A, Wong CF, Corliss HL, Lankenau SE. Risk factors for high levels of prescription drug misuse and illicit drug use among substance-using young men who have sex with men \(YMSM\). Drug Alcohol Depend 2015; 150:156.](#)
129. [Newcomb ME, Ryan DT, Greene GJ, et al. Prevalence and patterns of smoking, alcohol use, and illicit drug use in young men who have sex with men. Drug Alcohol Depend 2014; 141:65.](#)
130. [Halkitis PN, Siconolfi DE, Stults CB, et al. Modeling substance use in emerging adult gay, bisexual, and other YMSM across time: the P18 cohort study. Drug Alcohol Depend 2014; 145:209.](#)

131. [Rosario M, Reisner SL, Corliss HL, et al. Sexual-orientation disparities in substance use in emerging adults: a function of stress and attachment paradigms. Psychol Addict Behav 2014; 28:790.](#)
132. [Dai H. Tobacco Product Use Among Lesbian, Gay, and Bisexual Adolescents. Pediatrics 2017.](#)
133. [Phillips Ii G, Turner B, Felt D, et al. Trends in Alcohol Use Behaviors by Sexual Identity and Behavior Among High School Students, 2007-2017. J Adolesc Health 2019; 65:760.](#)
134. [Austin SB, Ziyadeh N, Fisher LB, et al. Sexual orientation and tobacco use in a cohort study of US adolescent girls and boys. Arch Pediatr Adolesc Med 2004; 158:317.](#)
135. [Corliss HL, Rosario M, Wypij D, et al. Sexual orientation disparities in longitudinal alcohol use patterns among adolescents: findings from the Growing Up Today Study. Arch Pediatr Adolesc Med 2008; 162:1071.](#)
136. [Russell ST, Driscoll AK, Truong N. Adolescent same-sex romantic attractions and relationships: implications for substance use and abuse. Am J Public Health 2002; 92:198.](#)
137. [Storholm ED, Halkitis PN, Siconolfi DE, Moeller RW. Cigarette smoking as part of a syndemic among young men who have sex with men ages 13-29 in New York City. J Urban Health 2011; 88:663.](#)
138. [Freeman P, Walker BC, Harris DR, et al. Methamphetamine use and risk for HIV among young men who have sex with men in 8 US cities. Arch Pediatr Adolesc Med 2011; 165:736.](#)
139. [Newcomb ME, Mustanski B. Developmental change in the relationship between alcohol and drug use before sex and sexual risk behavior in young men who have sex with men. AIDS Behav 2014; 18:1981.](#)
140. [Gambon TB, Gewirtz O'Brien JR, COMMITTEE ON PSYCHOSOCIAL ASPECTS OF CHILD AND FAMILY HEALTH, COUNCIL ON COMMUNITY PEDIATRICS. Runaway Youth: Caring for the Nation's Largest Segment of Missing Children. Pediatrics 2020; 145.](#)
141. [Noell JW, Ochs LM. Relationship of sexual orientation to substance use, suicidal ideation, suicide attempts, and other factors in a population of homeless adolescents. J Adolesc Health 2001; 29:31.](#)
142. [Kruks G. Gay and lesbian homeless/street youth: special issues and concerns. J Adolesc Health 1991; 12:515.](#)

143. [Corliss HL, Goodenow CS, Nichols L, Austin SB. High burden of homelessness among sexual-minority adolescents: findings from a representative Massachusetts high school sample. Am J Public Health 2011; 101:1683.](#)
144. [Rice E, Barman-Adhikari A, Rhoades H, et al. Homelessness experiences, sexual orientation, and sexual risk taking among high school students in Los Angeles. J Adolesc Health 2013; 52:773.](#)
145. [Cochran BN, Stewart AJ, Ginzler JA, Cauce AM. Challenges faced by homeless sexual minorities: comparison of gay, lesbian, bisexual, and transgender homeless adolescents with their heterosexual counterparts. Am J Public Health 2002; 92:773.](#)
146. [Whitbeck LB, Chen X, Hoyt DR, et al. Mental disorder, subsistence strategies, and victimization among gay, lesbian, and bisexual homeless and runaway adolescents. J Sex Res 2004; 41:329.](#)
147. [Van Leeuwen JM, Boyle S, Salomonsen-Sautel S, et al. Lesbian, gay, and bisexual homeless youth: an eight-city public health perspective. Child Welfare 2006; 85:151.](#)
148. [Marshall BD, Shannon K, Kerr T, et al. Survival sex work and increased HIV risk among sexual minority street-involved youth. J Acquir Immune Defic Syndr 2010; 53:661.](#)
149. [Weber AE, Boivin JF, Blais L, et al. Predictors of initiation into prostitution among female street youths. J Urban Health 2004; 81:584.](#)
150. [Weber AE, Boivin JF, Blais L, et al. HIV risk profile and prostitution among female street youths. J Urban Health 2002; 79:525.](#)
151. [Zou H, Prestage G, Fairley CK, et al. Sexual behaviors and risk for sexually transmitted infections among teenage men who have sex with men. J Adolesc Health 2014; 55:247.](#)
152. [Stults CB, Javdani S, Greenbaum CA, et al. Intimate partner violence and substance use risk among young men who have sex with men: The P18 cohort study. Drug Alcohol Depend 2015; 154:54.](#)
153. [Stults CB, Javdani S, Greenbaum CA, et al. Intimate Partner Violence and Sex Among Young Men Who Have Sex With Men. J Adolesc Health 2016; 58:215.](#)
154. [Ybarra ML, Rosario M, Saewyc E, Goodenow C. Sexual Behaviors and Partner Characteristics by Sexual Identity Among Adolescent Girls. J Adolesc Health 2016; 58:310.](#)
155. [Perry NS, Nelson KM, Carey MP. Diversity of Psychosocial Syndemic Indicators and](#)

- [Associations with Sexual Behavior with Male and Female Partners Among Early Adolescent Sexual Minority Males. *LGBT Health* 2019; 6:386.](#)
156. [Herrick AL, Marshal MP, Smith HA, et al. Sex while intoxicated: a meta-analysis comparing heterosexual and sexual minority youth. *J Adolesc Health* 2011; 48:306.](#)
157. [Centers for Disease Control and Prevention \(CDC\). Vital signs: HIV infection, testing, and risk behaviors among youths - United States. *MMWR Morb Mortal Wkly Rep* 2012; 61:971.](#)
158. Centers for Disease Control and Prevention. HIV Surveillance Report, 2012; vol. 24. Published November 2014. Accessed October 2016. www.cdc.gov/hiv/library/reports/surveillance/.
159. CDC. HIV surveillance in adolescents and young adults. <http://www.cdc.gov/hiv/topics/surveillance/resources/slides/adolescents/index.htm> (Accessed on June 13, 2011).
160. [Halkitis P, Kapadia F, Ompad D. Incidence of HIV Infection in Young Gay, Bisexual, and Other YMSM: The P18 Cohort Study. *J Acquir Immune Defic Syndr* 2015; 69:466.](#)
161. [Johnson AS, Hall HI, Hu X, et al. Trends in diagnoses of HIV infection in the United States, 2002-2011. *JAMA* 2014; 312:432.](#)
162. [Koh AS, Gómez CA, Shade S, Rowley E. Sexual risk factors among self-identified lesbians, bisexual women, and heterosexual women accessing primary care settings. *Sex Transm Dis* 2005; 32:563.](#)
163. [Kwakwa HA, Ghobrial MW. Female-to-female transmission of human immunodeficiency virus. *Clin Infect Dis* 2003; 36:e40.](#)
164. [Chan SK, Thornton LR, Chronister KJ, et al. Likely female-to-female sexual transmission of HIV--Texas, 2012. *MMWR Morb Mortal Wkly Rep* 2014; 63:209.](#)
165. [Committee Opinion No 708: Improving Awareness of and Screening for Health Risks Among Sex Workers. *Obstet Gynecol* 2017; 130:e53.](#)
166. [Saewyc EM, Bearinger LH, Blum RW, Resnick MD. Sexual intercourse, abuse and pregnancy among adolescent women: does sexual orientation make a difference? *Fam Plann Perspect* 1999; 31:127.](#)
167. [Saewyc EM, Poon CS, Honna Y, Skay CL. Stigma management? The links between enacted stigma and teen pregnancy trends among gay, lesbian, and bisexual students in British Columbia. *Can J Hum Sex* 2008; 17:123.](#)
168. [Herrick AL, Matthews AK, Garofalo R. Health risk behaviors in an urban sample of](#)

- [young women who have sex with women. J Lesbian Stud 2010; 14:80.](#)
169. [Charlton BM, Nava-Coulter B, Coles MS, Katz-Wise SL. Teen Pregnancy Experiences of Sexual Minority Women. J Pediatr Adolesc Gynecol 2019; 32:499.](#)
170. [Blake SM, Ledsky R, Lehman T, et al. Preventing sexual risk behaviors among gay, lesbian, and bisexual adolescents: the benefits of gay-sensitive HIV instruction in schools. Am J Public Health 2001; 91:940.](#)
171. [Lindley LL, Walsemann KM. Sexual Orientation and Risk of Pregnancy Among New York City High-School Students. Am J Public Health 2015; 105:1379.](#)
172. [Goldberg SK, Reese BM, Halpern CT. Teen Pregnancy Among Sexual Minority Women: Results From the National Longitudinal Study of Adolescent to Adult Health. J Adolesc Health 2016; 59:429.](#)
173. [Charlton BM, Roberts AL, Rosario M, et al. Teen Pregnancy Risk Factors Among Young Women of Diverse Sexual Orientations. Pediatrics 2018; 141.](#)
174. [Saewyc EM, Pettingell S, Magee LL. The prevalence of sexual abuse among adolescents in school. J Sch Nurs 2003; 19:266.](#)
175. [Blashill AJ, Safren SA. Sexual orientation and anabolic-androgenic steroids in U.S. adolescent boys. Pediatrics 2014; 133:469.](#)
176. [Matthews-Ewald MR, Zullig KJ, Ward RM. Sexual orientation and disordered eating behaviors among self-identified male and female college students. Eat Behav 2014; 15:441.](#)
177. [Robin L, Brener ND, Donahue SF, et al. Associations between health risk behaviors and opposite-, same-, and both-sex sexual partners in representative samples of vermont and massachusetts high school students. Arch Pediatr Adolesc Med 2002; 156:349.](#)
178. [Austin SB, Ziyadeh NJ, Corliss HL, et al. Sexual orientation disparities in purging and binge eating from early to late adolescence. J Adolesc Health 2009; 45:238.](#)
179. [Ackard DM, Fedio G, Neumark-Sztainer D, Britt HR. Factors associated with disordered eating among sexually active adolescent males: gender and number of sexual partners. Psychosom Med 2008; 70:232.](#)
180. [Austin SB, Ziyadeh N, Kahn JA, et al. Sexual orientation, weight concerns, and eating-disordered behaviors in adolescent girls and boys. J Am Acad Child Adolesc Psychiatry 2004; 43:1115.](#)

181. [Hadland SE, Austin SB, Goodenow CS, Calzo JP. Weight misperception and unhealthy weight control behaviors among sexual minorities in the general adolescent population. J Adolesc Health 2014; 54:296.](#)
182. [Katz-Wise SL, Blood EA, Milliren CE, et al. Sexual orientation disparities in BMI among U.S. adolescents and young adults in three race/ethnicity groups. J Obes 2014; 2014:537242.](#)
183. [Reisner SL, Van Wagenen A, Gordon A, Calzo JP. Disparities in safety belt use by sexual orientation identity among US high school students. Am J Public Health 2014; 104:311.](#)
184. [Needham BL, Austin EL. Sexual orientation, parental support, and health during the transition to young adulthood. J Youth Adolesc 2010; 39:1189.](#)
185. [Puckett JA, Woodward EN, Mereish EH, Pantalone DW. Parental Rejection Following Sexual Orientation Disclosure: Impact on Internalized Homophobia, Social Support, and Mental Health. LGBT Health 2015; 2:265.](#)
186. [Huebner DM, Roche KM, Rith KA. Effects of Family Demographics and the Passage of Time on Parents' Difficulty with Their Lesbian, Gay, or Bisexual Youth's Sexual Orientation. Arch Sex Behav 2019; 48:1581.](#)
187. Report of the American Psychological Association Task Force on Appropriate Therapeutic Responses to Sexual Orientation. 2009 <http://www.apa.org/pi/lgbt/resources/therapeutic-response.pdf> (Accessed on January 23, 2014).
188. [APA. Commission on Psychotherapy by Psychiatrists.. Position statement on therapies focused on attempts to change sexual orientation \(reparative or conversion therapies\). Am J Psychiatry 2000; 157:1719.](#)
189. [Rossi NE. "Coming out" stories of gay and lesbian young adults. J Homosex 2010; 57:1174.](#)
190. [D'Augelli AR, Hershberger SL, Pilkington NW. Lesbian, gay, and bisexual youth and their families: disclosure of sexual orientation and its consequences. Am J Orthopsychiatry 1998; 68:361.](#)
191. [Kosciw JG, Palmer NA, Kull RM. Reflecting resiliency: openness about sexual orientation and/or gender identity and its relationship to well-being and educational outcomes for LGBT students. Am J Community Psychol 2015; 55:167.](#)
192. [McLaughlin KA, Hatzenbuehler ML, Xuan Z, Conron KJ. Disproportionate exposure to](#)

[early-life adversity and sexual orientation disparities in psychiatric morbidity. Child Abuse Negl 2012; 36:645.](#)


193. Higgitt N, Wingert S, Ristock J, et al. Voices from the Margins: Experiences of Street-Involved Youth in Winnipeg. University of Winnipeg, 2003. Available at: www.uwinnipeg.ca/faculty/ius/iusweb_backup/pdf/Street-kidsReportfinalSeptember903.pdf (Accessed on May 30, 2013).
194. Biegel S, Kuehl SJ. Safe at School: Addressing the School Environment and LGBT Safety through Policy and Legislation. National Education Policy Center October 2010. Available at: www.scribd.com/doc/48054580/Biegel-LGBT#archive (Accessed on May 30, 2013).
195. [Birkett M, Espelage DL, Koenig B. LGB and questioning students in schools: the moderating effects of homophobic bullying and school climate on negative outcomes. J Youth Adolesc 2009; 38:989.](#)
196. [McGuire JK, Anderson CR, Toomey RB, Russell ST. School climate for transgender youth: a mixed method investigation of student experiences and school responses. J Youth Adolesc 2010; 39:1175.](#)
197. [De Pedro KT, Esqueda MC, Gilreath TD. School Protective Factors and Substance Use Among Lesbian, Gay, and Bisexual Adolescents in California Public Schools. LGBT Health 2017; 4:210.](#)
198. [Russell ST, Ryan C, Toomey RB, et al. Lesbian, gay, bisexual, and transgender adolescent school victimization: implications for young adult health and adjustment. J Sch Health 2011; 81:223.](#)
199. Russell ST, McGuire JK. The school climate for lesbian, gay, bisexual, and transgender (LGBT) students. In: Toward Positive Youth Development: Transforming Schools and Community Programs, Shinn M, Yoshikawa H (Eds), Oxford University Press, Oxford, UK 2008. p.133.
200. O'Shaughnessy M, Russell ST, Heck K, et al. Safe Place to Learn: Consequences of Harassment Based on Actual or Perceived Sexual Orientation and Gender Non-Conformity and Steps for Making Schools Safer. California Safe Schools Coalition, 2004. www.casafeschools.org/SafePlacetoLearnLow.pdf (Accessed on January 29, 2014).
201. [Szalacha L. Safe sexual diversity climates: lessons learned from an evaluation of Massachusetts Safe Schools Program for Gay and Lesbian Students. Am J Educ 2003; 110:58.](#)

202. [Toomey RB, Ryan C, Diaz RM, et al. Gender-nonconforming lesbian, gay, bisexual, and transgender youth: school victimization and young adult psychosocial adjustment. Dev Psychol 2010; 46:1580.](#)
203. [Proulx CN, Coulter RWS, Egan JE, et al. Associations of Lesbian, Gay, Bisexual, Transgender, and Questioning-Inclusive Sex Education With Mental Health Outcomes and School-Based Victimization in U.S. High School Students. J Adolesc Health 2019; 64:608.](#)

Topic 90691 Version 44.0

GRAPHICS


Paradigm of sexuality



Birth-assigned gender - Brain, hormones, body parts assigning male/female gender, usually at birth

Gender identity - Person's basic sense of being male, female, both, neither, or something else entirely – especially as experienced in self-awareness and behavior

Gender expression - Ways in which person acts, presents self, and communicates gender within a given culture



LGBTQQI - Lesbian, Gay, Bisexual, Transgender, Queer, Questioning, Intersex

YMSM - Young Men who have Sex with Men


YWSW - Young Women who have Sex with Women

Bisexual, pansexual, asexual, queer

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 50053 Version 7.0

Minority stress theory



Sexual minoritized youth (eg, lesbian, gay, bisexual, transgender) may experience stigma related to their sexual orientation or gender diversity. This stigma can lead to prejudice, discrimination, and lack of acceptance in society at large or in one's relationships with peers and family members, resulting in "minority stress." Stigmatization and minority stress may increase vulnerability to psychologic distress (eg, anxiety, depression), which may contribute to adverse health and psychosocial outcomes (eg, suicide, substance use, socioeconomic disadvantage, victimization). Psychologic distress and adverse psychosocial outcomes are society-induced and not inherent to being a sexual minoritized individual.

SES: socioeconomic status.

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95318 Version 5.0

Strategies for interviewing parents about their child's sexual attraction, identity, and behaviors

Sample questions regarding sexuality	Developmental goals
For parent(s) of prepubertal or peripubertal adolescents (age <14 years)	
<ul style="list-style-type: none"> ■ Do you have any questions or concerns about how your child expresses gender or sexuality? ■ What sorts of play, interests, and friends does your child have? ■ Does any of your child's play, activities, or interests concern you or your co-parent? ■ Do you have any concerns about your child's friends, lack of friends, bullying, or other social issues? ■ Do you need information regarding upcoming changes with puberty and adolescence? ■ How do you plan to talk with your child about puberty? ■ What were your experiences with puberty, adolescence, and sex? ■ What sort of expectations and plan for communication have you established with your teen? 	<ul style="list-style-type: none"> ■ Preparing parents to tasks of puberty ■ Preparing parents to talk with children about upcoming puberty changes ■ Prepare parents to have open, honest, supportive communication ■ Give parents skillset to set limits, establish plans, and discuss consequences

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95292 Version 3.0

Strategies for interviewing prepubertal and peripubertal adolescents about sexual attraction, identity, and behaviors

Sample questions regarding sexuality	Developmental goals
For prepubertal children and peripubertal adolescents (age <10 years)	
<ul style="list-style-type: none"> Do you have any questions or concerns about what being a boy or girl means? How do you feel – more like a boy, girl, or somewhere in between? Do you like your body? Do you think your body is healthy? Does your body fit you? What do you wish was different about your body? If you could change anything about your body, what would it be? Have you talked with your parent(s) or at school about puberty? What does puberty mean to you? Do you know what changes are ahead when you start puberty? What sort of plan do you have with your parent(s) for talking about teen issues? 	<ul style="list-style-type: none"> Prepare children to enter puberty Prepare children to have open, honest, supportive communication Give children permission to ask questions Give children a skillset to ask questions Give children an explicit plan about how to bring up sensitive topics with adults
For peripubertal and early adolescents (age 10 to 14 years)	
<ul style="list-style-type: none"> What do you know about puberty? What sort of conversations have you had about being a teen with your parents, at school, or with friends? How do you feel about the changes in your body? Have your friends' bodies started to change? What changes are occurring with your friends regarding crushes, romantic feelings, dating, or sex? Do you have any crushes or romantic interests in your life? Is there anyone who is extra special and who you would like to get to know or date? Most kids masturbate. Do you get aroused, touch yourself, or wonder about having sex? Some kids start experimenting with sex early. Do you have any questions about what sex is? What are your friends doing in regards to sex? What have you done so far about sex? What and how do you and your parent(s) communicate about sex? What is your parent(s) position on being sexually active? Can you talk openly with your parent(s) about sex? Is there anything I can do to help you communicate with your parent about puberty and sex? 	<ul style="list-style-type: none"> Early education, preparation for puberty Early identification of gender dysphoria, same-sex attraction and behaviors Early support for sexual minoritized youth and early intervention for parents (eg, resources, referrals) Risk reduction and prevention of: anxiety, depression, suicidality, and other mood or behavioral concerns

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95294 Version 3.0

Strategies for interviewing middle and late adolescents about sexual attraction, identity, and behaviors

Sample questions regarding sexuality	Developmental goals
For middle adolescents (age 15 to 17 years)	
<ul style="list-style-type: none"> Do you have any romantic interests in your life? Are you dating anyone? Is there anyone with whom you have a sexual relationship, including kissing, touching, or other stuff? Tell me about your partner. Are you sexually active or have you ever had sex*? What do you mean by "sex"? What sorts of sex have you had? What parts went where when you were having sex? Has sex been something you have wanted or something you felt emotionally or physically forced into doing? Have you ever paid for sex or been paid for sex? Who have you talked with about sex? What do your parents have to say about sex? What do your parents, family, and friends say about same-sex or gay people? Do you have any gay or bisexual people in your life that you can talk with? How do you feel about homosexuality or same-sex relationships? 	<ul style="list-style-type: none"> Safer exploration, experimentation Risk reduction and prevention of: sexually transmitted infections, unintended pregnancy, dating violence, bullying, substance use, mental health concerns, and suicidality Acknowledging and modeling support for diversity
For late adolescents (age 18 and older)	
<ul style="list-style-type: none"> Are you sexually active or have you ever had sex*? Why or why not? What benefits or risks does being sexually active bring to a relationship? What sorts of things do you and your partner(s) do when you are sexually active? Do you kiss, touch each other's genitals, put your fingers in your partner's vagina or butt, put your penis in your partner's vagina or butt, receive a penis in your vagina or butt, give or receive oral sex? Has sex been something you have wanted or something you felt emotionally or physically forced into doing? Have you ever paid for sex or been paid for sex? Is sex pleasurable for you? Do you have any concerns about sex, orgasm? Any concerns about premature ejaculation or having a hard time getting aroused, wet, or hard? Do you do anything else that gives you and/or your partner sexual pleasure? Masturbation? Pornography? Drugs or alcohol with sex? Sex games or fantasy? Sadism or masochism? Fetishes or unusual ways to pleasure yourself or partner? 	<ul style="list-style-type: none"> Risk and harm reduction Explore sex for pleasure and pleasurable sex Explore sex's role in intimacy and mature relationship building Prepare for intimate partners, family planning

- (Remember to **leave long and open pauses** for patients to fill in the blanks or report other activities or interests)

* For those who are sexually active, it is also important to ask about the number of partners, use of condoms or dental dams, and contraception.

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95293 Version 3.0

Resources for parents and family members of sexual minoritized youth

<p>Advocates for Youth: LGBTQ Issues Info for Parents</p> <p>https://advocatesforyouth.org/resources-tools/?_sft_type=health-information&_sft_audience=for-parents</p>
<p>American Academy of Pediatrics</p> <p>Gay, lesbian, bisexual teens: Facts for teens and their parents</p> <p>www.healthychildren.org/English/ages-stages/teen/dating-sex/Pages/Gay-Lesbian-and-Bisexual-Teens-Facts-for-Teens-and-Their-Parents.aspx</p>
<p>American Academy of Child and Adolescent Psychiatry</p> <p>Facts for Families: Gay, Lesbian and Bisexual Adolescents</p> <p>https://www.aacap.org/AACAP/Families_and_Youth/Facts_for_Families/FFF-Guide/Gay-Lesbian-and-Bisexual-Adolescents-063.aspx</p>
<p>American Psychological Association</p> <p>Understanding sexual orientation and gender identity</p> <p>www.apa.org/topics/sexuality/index.aspx</p>
<p>Centers for Disease Control and Prevention</p> <p>Electronic aggression: Technology and youth violence</p> <p>https://www.cdc.gov/violenceprevention/pdf/ea-tipsheet-a.pdf</p>
<p>Kids Health: Sexual Attraction and Orientation</p> <p>kidshealth.org/parent/emotions/feelings/sexual_orientation.html</p>
<p>Parents, Families, and Friends of Lesbians and Gays</p> <p>pflag.org/Family</p>
<p>Stopbullying.gov</p> <p>www.stopbullying.gov</p>
<p>Stop Bullying Now</p> <p>www.stopbullyingnow.com</p>
<p>Supportive Families, Healthy Children: Helping Families with Lesbian, Gay, Bisexual and Transgender Children</p> <p>familyproject.sfsu.edu/sites/sites7.sfsu.edu.familyproject/files/FAP_English%20Booklet_pst.pdf</p>
<p>The Trevor Project: Education and training for adults</p> <p>www.thetrevorproject.org/section/education-training-for-adults</p>

LGBTQ: lesbian, gay, bisexual, transgender, questioning.

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95287 Version 11.0

Discussing disclosure with sexual minoritized youth

When?	<p>Coming out can be a lifelong process. Everyone's timing is different. The person coming out should decide when is best.</p> <p>Consider coming out when you:</p> <ul style="list-style-type: none"> ▪ Are comfortable with your sexual interests, identity, and orientation ▪ Want to share this with persons that you trust and are close to you ▪ Have had a chance to gather information or talk to others to learn from their coming out experiences ▪ Have a plan for support – where could you live, get money for food and clothes, go to school, and work if you are not accepted in your home and need to live elsewhere? ▪ Are ready to
Who?	<p>You don't have to tell everyone at the same time. Parents and family may be a strong source of support but also may respond differently than you expect. Sometimes telling a close and trusted friend or professional (eg, teacher, coach, counselor) can be easier and safer the first time you tell someone about your sexuality.</p> <p>Choose a person or persons who you:</p> <ul style="list-style-type: none"> ▪ Know well and expect to be respectful and supportive ▪ Trust, feel safe with, and know will keep your information confidential if you ask them to ▪ If you are thinking about telling a friend, choose carefully; some may have trouble keeping your information safe and confidential ▪ If you are thinking about disclosing to a school staff member, it is important to know whether or not your school's privacy and confidentiality policy requires staff to reveal that information to your parents ▪ Be sure to tell the person or persons you tell, who else may be told and who should not be told
How?	<p>Planning</p> <p>Planning for your disclosure can help to make it a more positive and safe experience. Preparation may include:</p> <ul style="list-style-type: none"> ▪ Talking to other sexual minoritized youth or sexual minoritized adults who have already come out to family and friends; listening and learning from others may provide useful strategies and help you feel confident about when and how to come out ▪ Participating in LGBTQ youth groups, support groups, or school groups that can offer advice on coming out ▪ Accessing internet resources that relate other youths' coming out experiences ▪ Creating a support network before you come out; professionals (eg, health care providers, mental health professionals, counselors) who work with children and teens may be able to help with this <p>Process</p> <p>Disclosing in a letter</p> <p>Writing a letter gives you time to carefully say what you mean to say. Reading a letter in private gives the other person time to consider how they feel and how they might respond to you.</p> <p>Disclosing in person</p> <p>Some things to keep in mind if you decide to disclose in person:</p> <ul style="list-style-type: none"> ▪ Try to choose a quiet and private space where you can talk with one or two people at a time. ▪ If you have concerns about your parents' reaction (or the reaction of whoever you are telling), try to choose a quiet but public place with other persons around and available to help if needed. ▪ Disclose when you and the other person are well rested, not dealing with other active stressors, and have time to continue talking. ▪ Focus on a few issues that are most important to you. Actively listen to what is important to the other person as well.

	<ul style="list-style-type: none"> Plan how you might want to end the conversation and how you might plan to talk about it later. Avoid using alcohol or drugs; your mental and emotional state should be stable and secure when you have this important conversation. Try to avoid coming out because of pressure from others, being angry, or fighting.
Safety	Be prepared for the possibility of a negative or rejecting reaction.
	Safety first and foremost. Plan how to keep yourself physically safe at the time of disclosure and afterward. This may be difficult, but it is important to figure out ahead of time where you would go if the disclosure is difficult or threatening.
	Emotional safety is important, too. If you think you may be disclosing to a person who will be negative or rejecting, have your support network ready to help you process the negative reaction.
	Some people's first reaction may be more negative than you anticipate. The negative response may be based on the initial shock, surprise, or discomfort. They may need time to process the disclosure. People with negative initial reactions may still accept and support you but have a hard time processing their own feelings.
	Strong negative reactions may reflect personal biases or a background and culture that discriminate against sexual minorities. Issues of religion, morality, and culture have a large impact on how people view sexual diversity. However, there are many social and religious institutions that support civil and social rights of individuals regardless of sexuality or orientation.
	You get to decide if and when to walk away from people who are negative, unsupportive, or harmful.

LGBTQ: lesbian, gay, bisexual, transgender, questioning.

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95291 Version 3.0

Resources for sexual minoritized youth and their friends

Gay-Straight Alliance Network gsanetwork.org
Gay Lesbian Straight Education Network: Information for Students glSEN.org/students
It Gets Better Project www.itgetsbetter.org
Sexuality Information and Education Council of the United States www.siecus.org
Stopbullying.gov www.stopbullying.gov
Stop Bullying Now www.stopbullyingnow.com
The Trevor Project: Help and Suicide Prevention www.thetrevorproject.org
Trevor Space: News and Networking www.trevorspace.org

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95288 Version 4.0

Health-related national organizations/coalitions for sexual minoritized persons

American Psychiatric Association - LGBT Sexual Orientation www.psychiatry.org/mental-health/people/lgbt-sexual-orientation
Gay and Lesbian Medical Association www.glma.org
National Association of Lesbian and Gay Addiction Professionals www.nalgap.org
National Coalition for LGBT Health http://healthlgbt.org/
National LGBT Cancer Network www.cancer-network.org
National LGBT Health Education Center www.lgbthealtheducation.org
National LGBT Tobacco Control Network www.lgbttobacco.org
Society for Adolescent Health and Medicine Position Statement on LGBT Adolescent Health and Well-being www.adolescenthealth.org/Advocacy/Position-Papers-Statements.aspx

LGBT: lesbian, gay, bisexual, transgender.

Courtesy of Michelle Forcier, MD, MPH and Johanna Olson-Kennedy, MD.

Graphic 95289 Version 8.0

Contributor Disclosures

Michelle Forcier, MD, MPH Consultant/Advisory Boards: Planned Parenthood. Other Financial Interest: Springer [Gender book royalties]. **Johanna Olson-Kennedy, MD** Consultant/Advisory Boards: Novartis [Transgender adolescents and young adults]. Consultant/Advisory Boards (Spouse/partner): Novartis [Transgender adolescents and young adults]. **Diane Blake, MD** Nothing to disclose **Mary M Torchia, MD** Nothing to disclose

Contributor disclosures are reviewed for conflicts of interest by the editorial group. When found, these are addressed by vetting through a multi-level review process, and through requirements for references to be provided to support the content. Appropriately referenced content is required of all authors and must conform to UpToDate standards of evidence.

[Conflict of interest policy](#)

→